

胞核因子 β 悅洗飄G上皮細胞核之表現 72 圖二十二、第四型肝細胞核因子 β 悅洗駢 β B膽及鰓之表現 73 圖二十三、第四型肝細胞核因子 β 悅洗鞠菱式B肌肉、睪丸及卵巢之表現 74 表目錄表一、選殖吳郭魚HNF-4 β 桶互q退化性引子之序列 76 表二、選殖吳郭魚HNF-4 β 掙廢搗M一性引子之序列 77 表三、選殖吳郭魚HNF-4 β 搗瑯搗M一性引子之序列 78 表四、HNF-4 β 桶刪T-PCR專一性引子 79 表五、吳郭魚HNF-4 β 宸穉]與其他物種間之序列比對 80 表六、各物種間第四型肝細胞核因子 β 琮皇i基酸序列之比對 81 表七、各物種間第四型肝細胞核因子 β 蚰皇i基酸序列之比對 82 附錄 附錄一、各物種間HNF-4 β 啞i基酸序列比對之相似度及相同度百分比 83 附錄二、各物種間HNF-4 β 啞i基酸DNA結合區之相似度及相同度百分比 84 附錄三、各物種間HNF-4 β 啞i基酸序列雙聚體/配位體結合區之相似度及相同度百分比 85

參考文獻

邵廣昭, 1996. 台灣常見魚貝類圖說(下). 台灣省漁業局. pp. 125-146. 王水良、房建中和傅繼梁。(2003). 肝富集轉錄因子及其相互作用研究進展. 國外醫學遺傳學分冊. 26(1):20-25. Argyrokastritis, A., Kamakari, S., Kapsetaki, M., Kritis, A., Talianidis, I. and Moschonas, N. K. (1997). Human hepatocyte nuclear factor-4 (hHNF-4) gene maps to 20q12-q13.1 between PLCG1 and D20S17. *Hum Genet*, 99(2), 233-236. Au-Fliegner, M., Helmer, E., Casasova, J., Raaka, B. and Samuels, H. (1993). The conserved ninth C-terminal heptad in thyroid hormone and retinoic acid receptors mediates diverse responses by affecting heterodimer but not homodimer formation. *Mol Cell Biol*, 13(9), 5725-5737. Bogan, A. A., Cohen, F. E. and Scanlan, T. S. (1998). Natural ligands of nuclear receptors have conserved volumes. *Nature Struct*, 5(10), 679-681. Bourguet, W., Vivat, V., Wurtz, J.-M., Chambon, P., Gronemeyer, H. and Moras, D. (2000). Crystal structure of a heterodimeric complex of RAR and RXR ligand-binding domains. *Mol Cell*, 5(2), 289-298. Carew, J. A., Pollak, E. S., Lopaciuk, S., and Bauer, K. A. (2000). A new mutation in the HNF4 binding region of the factor VII promoter in a patient with severe factor VII deficiency. *Blood*, 96(13), 4370-4372. Cereghini, S., Yaniv, M. and Cortese, R. (1990). Hepatocyte dedifferentiation and extinction is accompanied by a block in the synthesis of mRNA coding for the transcription factor HNF1/LFB1. *Embo J*, 9(7), 2257-2263. Chartier, F. L., Bossu, J. P., Laudet, V., Fruchart, J. C. and Laine, B. (1994). Cloning and sequencing of cDNAs encoding the human hepatocyte nuclear factor 4 indicate the presence of two isoforms in human liver. *Gene*, 147(2), 269-272. Chen, K. J., Chao, H. K., Hsiao, K. J., and Su, T. S. (2002). Identification and characterization of a novel liver-specific enhancer of the human phenylalanine hydroxylase gene. *Hum Genet*, 110(3), 235-243. Chen, M., Hieng, S., Qian, X., Costa, R. and Ou, J. H. (1994). Regulation of hepatitis B virus EN1 enhancer activity by hepatocyte-enriched transcription factor HNF3. *Virology*, 205(1), 127-132. Dhe-Paganon, S., Duda, K., Iwamoto, M., Chi, Y. I. and Shoelson, S. E. (2002). Crystal structure of the HNF4 alpha ligand binding domain in complex with endogenous fatty acid ligand. *J Biol Chem*, 277(41), 37973-37976. Drewes, T., Senkel, S., Holewa, B. and Ryffel, G. U. (1996). Human hepatocyte nuclear factor 4 isoforms are encoded by distinct and differentially expressed genes. *Mol Cell Biol*, 16(3), 925-931. Duncan, S. A., Manova, K., Chen, W. S., Hoodless, P., Weinstein, D. C., Bachvarova, R. F. (1994). Expression of transcription factor HNF-4 in the extraembryonic endoderm, gut, and nephrogenic tissue of the developing mouse embryo: HNF-4 is a marker for primary endoderm in the implanting blastocyst. *Proc Natl Acad Sci U S A*, 91(16), 7598-7602. Duncan, S. A., Nagy, A. and Chan, W. (1997). Murine gastrulation requires HNF-4 regulated gene expression in the visceral endoderm: tetraploid rescue of Hnf-4(-/-) embryos. *Development*, 124(2), 279-287. Egea, P. F., Mitschler, A., Rochel, N., Ruff, M., Chambon, P. and Moras, D. (2000). Crystal structure of the human RXR ligand-binding domain bound to its natural ligand: 9-cis retinoic acid. *EMBO J*, 19(11), 2592-2601. Forman, B. M., Yan, C.-R., Au, M., Casanova, J., Ghysdal, J. and Samuels, H. H. (1998). A domain containing leucine-zipper-like motifs mediate novel in vivo interactions between the thyroid hormone and retinoic acid receptors. *Mol Endocrinol*, 3(10), 1610-1626. Furuta, H., Iwasaki, N., Oda, N., Hinokio, Y., Horikawa, Y., Yamagata, K. (1997). Organization and partial sequence of the hepatocyte nuclear factor-4 alpha/MODY1 gene and identification of a missense mutation, R127W, in a Japanese family with MODY. *Diabetes*, 46(10), 1652-1657. Green, V. J., Kokkotou, E. and Ladias, J. A. (1998). Critical structural elements and multitarget protein interactions of the transcriptional activator AF-1 of hepatocyte nuclear factor 4. *J Biol Chem*, 273(45), 29950-29957. Griffo, G., Hamon-Benais, C., Angrand, P. O., Fox, M., West, L., Lecoq, O. (1993). HNF4 and HNF1 as well as a panel of hepatic functions are extinguished and reexpressed in parallel in chromosomally reduced rat hepatoma-human fibroblast hybrids. *J Cell Biol*, 121(4), 887-898. Hadzopoulou-Cladaras, M., Kistanova, E., Evagelopoulou, C., Zeng, S., Cladaras, C. and Ladias, J. A. (1997). Functional domains of the nuclear receptor hepatocyte nuclear factor 4. *J Biol Chem*, 272(1), 539-550. Hall, R. K., Sladek, F. M. and Granner, D. K. (1995). The orphan receptors COUP-TF and HNF-4 serve as accessory factors required for induction of phosphoenolpyruvate carboxykinase gene transcription by glucocorticoids. *Proc Natl Acad Sci U S A*, 92(2), 412-416. Harish, S., Khanam, T., Mani, S. and Rangarajan, P. (2001). Transcriptional activation by hepatocyte nuclear factor-4 in a cell-free system derived from rat liver nuclei. *Nucleic Acids Res*, 29(5), 1047-1053. Hata, S., Inoue, T., Kosuga, K., Nakashima, T., Tsukamoto, T. and Osumi, T. (1995). Identification of two splice isoforms of mRNA for mouse hepatocyte nuclear factor 4 (HNF-4). *Biochim Biophys Acta*, 1260(1), 55-61. Hata, S., Tsukamoto, T. and Osumi, T. (1992). A novel isoform of rat hepatocyte nuclear factor 4 (HNF-4). *Biochim Biophys Acta*, 1131(2), 211-213. Hattersley, A. T. (1998). Maturity-onset diabetes of the young: clinical heterogeneity explained by genetic heterogeneity. *Diabet Med*, 15(1), 15-24. Hayashi, Y., Wang, W., Ninomiya, T., Ohta, K. and Itoh, H. (1999). Liver enriched transcription factor and differentiation of hepatocellular carcinoma. *Mol. Pathol.* 52 : 19-24. Hayhurst, G. P., Lee, Y. H., Lambert, G., Ward, J. M. and Gonzalez, F. J. (2001). Hepatocyte nuclear factor 4alpha (nuclear receptor 2A1) is essential for maintenance of hepatic gene expression and lipid homeostasis. *Mol Cell Biol*, 21(4), 1393-1403. Hertz, R., Magenheimer, J., Berman, I. and Bar-Tana, J. (1998). Fatty acyl-CoA thioesters are ligands of hepatic nuclear factor-4alpha.

Nature, 392(6675), 512-516. Holewa, B., Zapp, D., Drewes, T., Senkel, S. and Ryffel, G. U. (1997). HNF4beta, a new gene of the HNF4 family with distinct activation and expression profiles in oogenesis and embryogenesis of *Xenopus laevis*. *Mol Cell Biol*, 17(2), 687-694. Huang, W. T., Gong, H. Y., Lin, C. J., Weng, C. F., Chen, M. H. and Wu, J. L. (2001). Hepatocyte nuclear factors-1alpha, -1beta, and -3beta expressed in the gonad of tilapia (*Oreochromis mossambicus*). *Biochem Biophys Res Commun*, 288(4), 833-840. Huang, W. T., Yu, H. C., Yang, H. C., Lin, C. J., Gong, H. Y., Weng, C. F. and Wu, J. L. (2004). Studies on the expression and localization of hepatocyte nuclear factor (HNFs) in gonads of tilapia (*Oreochromis mossambicus*). In "Proc. for Comparative Endocrinology", Nara, Japan. P-054. Imamura, K. and Tanaka, T. (1982). Pyruvate kinase isozymes from rat. *Methods Enzymol*, 90, 150-165. Jiang, G., Nepomuceno, L., Hopkins, K. and Sladek, F. M. (1995). Exclusive homodimerization of the orphan receptor hepatocyte nuclear factor 4 defines a new subclass of nuclear receptors. *Mol Cell Biol*, 15(9), 5131-5143. Ktistaki, E. and Talianidis, I. (1997). Chicken ovalbumin upstream promoter transcription factors act as auxiliary cofactors for hepatocyte nuclear factor 4 and enhance hepatic gene expression. *Mol Cell Biol*, 17(5), 2790-2797. Kuo, C. J., Conley, P. B., Chen, L., Sladek, F. M., Darnell, J. E., Jr. and Crabtree, G. R. (1992). A transcriptional hierarchy involved in mammalian cell-type specification. *Nature*, 355(6359), 457-461. Ladas, J. A., Hadzopoulou-Cladaras, M., Kardassis, D., Cardot, P., Cheng, J., Zannis, V. (1992). Transcriptional regulation of human apolipoprotein genes ApoB, ApoCIII, and ApoAII by members of the steroid hormone receptor superfamily HNF-4, ARP-1, EAR-2, and EAR-3. *J Biol Chem*, 267(22), 15849-15860. Lannoy, V. J., Decaux, J. F., Pierreux, C. E., Lemaigre, F. P., and Rousseau, G. G. (2002). Liver glucokinase gene expression is controlled by the onecut transcription factor hepatocyte nuclear factor-6. *Diabetologia*, 45(8), 1136-1141. Lee, J., Gulick, T. and Moore, D. (1992). Thyroid hormone receptor dimerization function maps to a conserved subregion of the ligand binding domain. *Mol Endocrinol*, 6(11), 1867-1873. Lees, J. A., Fawell, S. E., White, R. and Parker, M. G. (1990). A 22-amino-acid peptide restores DNA-binding activity to dimerization-defective mutants of the estrogen receptor. *Mol Cell Biol*, 10(10), 5529-5531. Leng, X., Blanco, J., Tsai, S. Y., Ozato, K., O'Malley, B. W. and Tsai, M. J. (1995). Mouse retinoid X receptor contains a separable ligand-binding and transactivation domain in its E region. *Mol Cell Biol*, 15(1), 255-263. Li, J., Ning, G. and Duncan, S. A. (2000). Mammalian hepatocyte differentiation requires the transcription factor HNF-4alpha. *Genes Dev*, 14(4), 464-474. Lindner, T., Gragnoli, C., Furuta, H., Cockburn, B. N., Petzold, C., Rietzsch, H. (1997). Hepatic function in a family with a nonsense mutation (R154X) in the hepatocyte nuclear factor-4alpha/MODY1 gene. *J Clin Invest*, 100(6), 1400-1405. Mangelsdorf, D. J., Thummel, C., Beato, M., Herrlich, P., Schutz, G., Umesono, K., Blumberg, B., Kastner, P., Mark, M., Chambon, P. and Evans, R. M. (1995). The nuclear receptor superfamily: the second decade. *Cell*, 83(6), 835-839. Mietus-Snyder, M., Sladek, F. M., Ginsburg, G. S., Kuo, C. F., Ladas, J. A., Darnell, J. E., Jr. (1992). Antagonism between apolipoprotein A1 regulatory protein 1, Ear3/COUP-TF, and hepatocyte nuclear factor 4 modulates apolipoprotein CIII gene expression in liver and intestinal cells. *Mol Cell Biol*, 12(4), 1708-1718. Miura, N. and Tanaka, K. (1993). Analysis of the rat hepatocyte nuclear factor (HNF) 1 gene promoter: synergistic activation by HNF4 and HNF1 proteins. *Nucleic Acids Res*, 21(16), 3731-3736. Morrissey, E. E., Tang, Z., Sigrist, K., Lu, M. M., Jiang, F., Ip, H. S. (1998). GATA6 regulates HNF4 and is required for differentiation of visceral endoderm in the mouse embryo. *Genes Dev*, 12(22), 3579-3590. Nakhei, H., Lingott, A., Lemm, I., and Ryffel, G. U. (1998). An alternative splice variant of the tissue specific transcription factor HNF4alpha predominates in undifferentiated murine cell types. *Nucleic Acids Res*, 26(2), 497-504. Navas, M. A., Vaisse, C., Boger, S., Heimesaat, M., Kollee, L. A. and Stoffel, M. (2000). The human HNF-3 genes: cloning, partial sequence and mutation screening in patients with impaired glucose homeostasis. *Hum Hered*, 50 : 370-81. Nishiyama, C., Hi, R., Osada, S. and Osumi, T. (1998). Functional interactions between nuclear receptors recognizing a common sequence element, the direct repeat motif spaced by one nucleotide (DR-1). *J Biochem (Tokyo)*, 123(6), 1174-1179. Nolten, L. A., Steenbergh, P. H. and Sussenbach, J. S. (1996). The hepatocyte nuclear factor 3beta stimulates the transcription of the human insulin-like growth factor I gene in a direct and indirect manner. *J Biol Chem*, 271(50), 31846-31854. Odom, D. T., Zizlsperger, N., Gordon, D. B., Bell, G. W., Rinaldi, N. J., Murray, H. L. (2004). Control of pancreas and liver gene expression by HNF transcription factors. *Science*, 303(5662), 1378-1381. Palamarchuk, A. Y., Kavsan, V. M., Sussenbach, J. S. and Holthuisen, P. E. (1999). The chum salmon IGF-II gene promoter is activated by hepatocyte nuclear factor 3beta. *FEBS Lett*, 446(2-3), 251-255. Peiler, G., Bockmann, B., Nakhei, H. and Ryffel, G. U. (2000). Inhibitor of the tissue-specific transcription factor HNF4, a potential regulator in early *Xenopus* development. *Mol Cell Biol*, 20(23), 8676-8683. Perlmann, T., Umesono, K., Rangarajan, P. N., Forman, B. M., and Evans, R. M. (1996). Two distinct dimerization interfaces differentially modulate target gene specificity of nuclear hormone receptors. *Mol Endocrinol*, 10(8), 958-966. Pogge v Strandmann, E., Senkel, S. and Ryffel, G. U. (2000). Ectopic pigmentation in *Xenopus* in response to DCoH/PCD, the cofactor of HNF1 transcription factor/pterin-4alpha-carbinolamine dehydratase. *Mech Dev*, 91(1-2), 53-60. Rosen, E., Beninghof, E. and Koenig, R. (1993). Dimerization interfaces of thyroid hormone, retinoic acid, vitamin D and retinoid X receptors. *J Biol Chem*, 268(16), 11534-11541. Ruse, M. D., Jr., Privalsky, M. L. and Sladek, F. M. (2002). Competitive cofactor recruitment by orphan receptor hepatocyte nuclear factor 4alpha1: modulation by the F domain. *Mol Cell Biol*, 22(6), 1626-1638. Schrem, H., Klempnauer, J. and Borlak, J. (2002). Liver-enriched transcription factors in liver function and development. Part I: the hepatocyte nuclear factor network and liver-specific gene expression. *Pharmacol Rev*, 54(1), 129-158. Shih, D. Q., Dansky, H. M., Fleisher, M., Assmann, G., Fajans, S. S. and Stoffel, M. (2000). Genotype/phenotype relationships in HNF-4alpha/MODY1: haploinsufficiency is associated with reduced apolipoprotein (AII), apolipoprotein (CIII), lipoprotein(a), and triglyceride levels. *Diabetes*, 49(5), 832-837. Sladek, F. M. (1994). Orphan receptor HNF-4 and liver-specific gene expression. *Receptor*, 4(1), 64. Sladek, F. M. and Seidel, S. D. (2000). Hepatocyte nuclear factor 4? In *Nuclear Receptors and Genetic Diseases* (Burris, T. P. and McCabe, E., eds), Academic Press, London. In the press. Sladek, F. M., Zhong, W. M., Lai, E. and Darnell, J. E., Jr. (1990). Liver-enriched transcription factor HNF-4 is a novel member of the steroid hormone receptor superfamily. *Genes Dev*, 4(12B), 2353-2365. Sladek, R. and Giguere, V. (2000). Orphan nuclear

receptors: an emerging family of metabolic regulators. *Adv Pharmacol*, 47, 23-87. Sourdive, D. J., Transy, C., Garbay, S. and Yaniv, M. (1997). The bifunctional DCOH protein binds to HNF1 independently of its 4-alpha-carbinolamine dehydratase activity. *Nucleic Acids Res*, 25(8), 1476-1484. Soutoglou, E., Katrakili, N. and Talianidis, I. (2000). Acetylation regulates transcription factor activity at multiple levels. *Mol Cell*, 5(4), 745-751. Stoffel, M. and Duncan, S. A. (1997). The maturity-onset diabetes of the young (MODY1) transcription factor HNF4alpha regulates expression of genes required for glucose transport and metabolism. *Proc Natl Acad Sci U S A*, 94(24), 13209-13214. Taraviras, S., Monaghan, A. P., Schutz, G. and Kelsey, G. (1994). Characterization of the mouse HNF-4 gene and its expression during mouse embryogenesis. *Mech Dev*, 48(2), 67-79. Tian, J. M. and Schibler, U. (1991). Tissue-specific expression of the gene encoding hepatocyte nuclear factor 1 may involve hepatocyte nuclear factor 4. *Genes Dev*, 5(12A), 2225-2234. Torres-Padilla, M. E., Fougere-Deschatrette, C. and Weiss, M. C. (2001). Expression of HNF4alpha isoforms in mouse liver development is regulated by sequential promoter usage and constitutive 3' end splicing. *Mech Dev*, 109(2), 183-193. Torres-Padilla, M. E., Sladek, F. M. and Weiss, M. C. (2002). Developmentally regulated N-terminal variants of the nuclear receptor hepatocyte nuclear factor 4alpha mediate multiple interactions through coactivator and corepressor-histone deacetylase complexes. *J Biol Chem*, 277(47), 44677-44687. Torres-Padilla, M. E. and Weiss, M. C. (2003). Effects of interactions of hepatocyte nuclear factor 4alpha isoforms with coactivators and corepressors are promoter-specific. *FEBS Lett*, 539(1-3), 19-23. Whitfield, G. K., Hsieh, J. C., Nakajima, S., Macdonald, P. N., Thompson, P. D., Jurutka, P. W., Haussler, C. A. and Haussler, M. R. (1995). A highly conserved region in the hormone-binding domain of the human vitamin D receptor contains residues vital for transcriptional activation. *Mol Endocrinol*, 9(9), 1166-1179. Weber, A., Marie, J., Cottreau, D., Simon, M. P., Besmond, C., Dreyfus, J. C. and Kahn, A. (1984). Dietary control of aldolase B and L-type pyruvate kinase mRNAs in rat. Study of translational activity and hybridization with cloned cDNA probes. *J Biol Chem*, 259(3), 1798-1802. Woolf, A. S. (2000). Diabetes, genes, and kidney development. *Kidney Int*, 57(3), 1202-1203. Yamagata, K., Furuta, H., Oda, N., Kaisaki, P. J., Menzel, S., Cox, N. J. (1996). Mutations in the hepatocyte nuclear factor-4alpha gene in maturity-onset diabetes of the young (MODY1). *Nature*, 384(6608), 458-460. Yanai, K., Hirota, K., Taniguchi-Yanai, K., Shigematsu, Y., Shimamoto, Y., Saito, T. (1999). Regulated expression of human angiotensinogen gene by hepatocyte nuclear factor 4 and chicken ovalbumin upstream promoter-transcription factor. *J Biol Chem*, 274(49), 34605-34612. Yoshida, E., Aratani, S., Itou, H., Miyagishi, M., Takiguchi, M., Osumu, T. (1997). Functional association between CBP and HNF4 in trans-activation. *Biochem Biophys Res Commun*, 241(3), 664-669. Zhong, W., Sladek, F. M. and Darnell, J. E., Jr. (1993). The expression pattern of a Drosophila homolog to the mouse transcription factor HNF-4 suggests a determinative role in gut formation. *Embo J*, 12(2), 537-544. Zhong, W., Mirkovitch, J. and Darnell, J. E., Jr. (1994). Tissue-specific regulation of mouse hepatocyte nuclear factor 4 expression. *Mol Cell Biol*, 14(11), 7276-7284.