

# Effect of Aqueous Extracts of Herbal Tea on Growth of Propionibacterium acnes

賴民峻、吳芳禎

E-mail: 9707201@mail.dyu.edu.tw

## ABSTRACT

Propionibacterium acnes are one of the major reasons for causing acne, such as lipase secreted from the human sebaceous follicle to decompose triglycerides of sebum and then to release acetic acid, propionic acid and butyric acid so that results in skin ulceration and inflammation. Gas Chromatography (GC) was employed in this research to analysis the content of acetic acid, propionic acid and butyric acid. The aims of this research were to investigate inhibit effect of P. acnes by 5 and 10 mg/mL extracts of 12 herbal tea. Extracts of herbal tea at 5 mg/mL could effectively control growth of P. acnes. Dry weight of microorganism from Lavandula officinalis, Melissa officinalis and Calendula officinalis were only 0.2-0.4 mg, which had great effect of inhibition. In terms of 10 mg/mL, dry weight of P. acnes from Calendula officinalis was  $18.61 \pm 1.33$  mg, obviously higher than that of control group (P

Keywords : Propionibacterium acnes, Extracts of herbal tea, Acetic acid, Propionic acid, Butyric acid

## Table of Contents

封面內頁 簽名頁 授權書iii 中文摘要iv 英文摘要vi 誌謝viii 目錄ix 圖目錄xii 表目錄xiii 第一章 緒言1 第二章 文獻回顧3 第一節 痘瘡3 一?成因3 二?痘瘡病原菌4 第二節 痘瘡的種類與併發症6 一?非發炎性8 二?發炎性8 三?併發症9 第三節 痘瘡的治療方法10 一?外用藥物12 二?口服藥物治療14 三、物理性治療15 第四節 天然植物在化妝品上的開發與應用18 一?天然植物萃取物之市場概況和發展潛力18 二?天然植物萃取物於化妝品之應用21 三、天然植物成份治療痘瘡之研究22 第三章 材料與方法25 第一節 材料25 一?花草茶25 二?菌株25 三?化學試藥26 四?儀器26 第二節 方法26 一?花草茶萃取物製備26 二?菌數之測定27 三?痘瘡丙酸桿菌之培養27 四?菌重之測定28 五?相對感應因子之測定28 六?回收率測定29 七?測定樣品有機酸含量29 八?統計分析29 第四章 結果與討論31 第一節 相對感應因子之測定31 第二節 痘瘡丙酸桿菌之菌重測定34 第三節 痘瘡丙酸桿菌產生乙酸含量分析38 一?添加5 mg/mL花草茶水萃物對痘瘡丙酸桿菌乙酸產量之影響38 二?添加10 mg/mL花草茶水萃物對痘瘡丙酸桿菌乙酸產量之影響38 三?探討添加花草茶水萃取物對痘瘡丙酸桿菌乙酸產量與乾菌重之關係39 第四節 痘瘡丙酸桿菌產生丙酸含量分析45 一?添加5 mg/mL花草茶水萃物對痘瘡丙酸桿菌丙酸產量之影響45 二?添加10 mg/mL花草茶水萃物對痘瘡丙酸桿菌丙酸產量之影響45 三?探討添加花草茶水萃取物對痘瘡丙酸桿菌丙酸產量與乾菌重之關係47 第五節 痘瘡丙酸桿菌產生丁酸含量分析53 一?添加5 mg/mL花草茶水萃物對痘瘡丙酸桿菌丁酸產量之影響53 二?添加10 mg/mL花草茶水萃物對痘瘡丙酸桿菌丁酸產量之影響53 三?探討添加花草茶水萃取物對痘瘡丙酸桿菌丁酸產量與乾菌重之關係55 第五章 結論60 參考文獻61 附錄66 圖目錄 圖一、發炎性痘瘡病害的發展7 圖二、痘瘡之治療順序11 圖三?比較不同萃取物添加量之痘瘡丙酸桿菌乾菌重37 圖四?比較痘瘡丙酸桿菌於不同萃取物添加量下之乙酸產量44 圖五?比較痘瘡丙酸桿菌於不同萃取物添加量下之丙酸產量52 圖六?比較痘瘡丙酸桿菌於不同萃取物添加量下之丁酸產量59 表目錄 表一?三種青春痘光照療法的比較17 表二?草本植物應用在皮膚保養品之作用19 表三?相對感應因子?滯留時間?偵測極限及回收率之測定值33 表四?痘瘡丙酸桿菌之乾菌重36 表五?添加花草茶水萃取物對痘瘡丙酸桿菌乙酸產量之影響41 表六?痘瘡丙酸桿菌乙酸產率42 表七、痘瘡丙酸桿菌乙酸產量對乾菌重之比值43 表八?添加花草茶水萃取物對痘瘡丙酸桿菌丙酸產量之影響49 表九?痘瘡丙酸桿菌丙酸產率50 表十、痘瘡丙酸桿菌丙酸產量對乾菌重之比值51 表十一?添加花草茶水萃取物對痘瘡丙酸桿菌丁酸產量之影響56 表十二?痘瘡丙酸桿菌丁酸產率57 表十三、痘瘡丙酸桿菌丁酸產量對乾菌重之比值58

## REFERENCES

1. 吳敏綺、邱琬婷、徐自菱和林文勝。2004。中西醫會診 - 青春痘。第9-83頁。書泉出版社，台北。
2. 洪偉章、李金枝和陳榮秀。2000。化妝品原料及功能。第64-65頁。藝軒圖書出版社。台北。
3. 徐雅芬、廖美智和羅淑慧。2005。天然萃取物應用在保養品、化妝品及醫藥產業之發展契機。第53-79頁。財團法人生物技術開發中心。台北。
4. 陳柏熹(譯)。1996。微生物診斷學。第445-449頁。合記圖書出版社。台北。
5. 陳豪勇(譯)。2002。最新醫用微生物學。第356-357頁。藝軒圖書出版社。台北。
6. 陳聰敏。1993。厭氧菌-感染與診療。第211-212頁。藝軒圖書出版社。台北。
7. 曾文楷。2002。中草藥對於痘瘡病原菌與黑色素生成的影響。靜宜大學應用化學研究所碩士論文。台中。
8. 曾銘儀。2004。彩色圖解皮膚學。第60-61頁。合記圖書出版社，台北。
9. 黃鶴群。2006a。台灣藥妝保養品品牌手冊。第14-180頁。合記圖書出版社。台北。
10. 黃鶴群。2006b。藥妝保養品與醫學美容諮詢手冊。第7-261頁。藝軒圖書出版社。台北。
11. 楊國藩。1997。化妝品技術師和醫師的皮膚科學。第80-81頁。國興出版社，新竹。
12. 楊麗珍。2003。皮膚疾病彩色圖譜。第91-97頁。合記圖書出版社，台北。
13. 趙昭明。2004。還我一張潔淨的臉。第50-51頁。大樹林出版社，台北。
14. 羅淑慧。2003。藥妝品之發展

趨勢。第54-145頁。財團法人生物技術開發中心，台北。15.Akerstedt, J. and Vollset, I. 1996. Malassezla Pachydermatis with special reference to canine skin disease. *The British Veterinary Journal* 152:269-281. 16.Ashkenazi, H., Malik, Z., Harth, Y. and Nitzan, Y. 2003. Eradication of Propionibacterium acnes by its endogenic porphyrins after illumination with high intensity blue light. *Immunology and Medical Microbiology* 35:17-24. 17.Bojar, R. A. and Holland, K. T. 2004. Acne and Propionibacterium acnes. *Clinics in Dermatology* 22:375-379. 18.Bunch, J. E. and Clark, T. J. 1997. Determination of volatile acids in tobacco, tea, and coffee using derivatization-purge and trap gas chromatography-selected ion monitoring mass spectrometry. *Journal of Chromatographic Science* 35:206-208. 19.Farrar, M. D. and Ingham, E. 2004. Acne: Inflammation. *Clinics in Dermatology* 22: 380 – 384. 20.Gibbs, B. F., Itiaba, K., Crawhall, C., Cooper, B. A. and Mamer, O. A. 1973. A rapid gas chromatographic method for the quantitation of volatile fatty acids in urine propionic acid excretion in vitamin B12 deficiency. *Journal of Chromatography* 81:65-69. 21.Guy, R., Green, M. and Kealey, T. 1996. Modeling of acne in vitro. *The Journal of Investigative Dermatology* 106: 176-182. 22.Higaki, S. 2003. Lipase inhibitors for the treatment of acne. *Journal of Molecular Catalysis B: Enzymatic* 22:377-384. 23.Higaki, S., Kitagawa, T., Kagoura, M., Morohashi, M. and Yamagishi, T. 2000. Relationship between Propionibacterium acnes biotypes and Jumi-haidoku-to. *The Journal of Dermatology* 27:635-638. 24.Higaki, S. and Morohashi, M. 2003. Propionibacterium acnes lipase in seborrheic dermatitis and other skin diseases and unsei-in. *Drugs Under Experimental & Clinical Rese* 4:157-159. 25.Higaki, S., Nakamura, M., Morohashi, M., Hasegawa, Y. and Yamagishi, T. 1996a. Anti-lipase activity of kampo formulations Coptidis Rhizoma and alkaloids against Propionibacterium acnes. *The Journal of Dermatology* 23:310-314. 26.Higaki, S., Nakamura, M., Morohashi, M., Hasegawa Y. and Yamagishi, T. 1996b. Activity of eleven kampo formulations and eight kampo crude drugs against Propionibacterium acnes isolated from acne patients: retrospective evaluation in 1990 and 1995. *The Journal of Dermatology* 23:871-875. 27.Higaki, S., Nakamura, M., Morohashi, M., and Yamagishi, T. 2004. Propionibacterium acnes biotypes and susceptibility to minocycline and Keigai-rengyo-to. *The Journal of Dermatology* 43:103-107. 28.Holland, K.T., Ingham, E. and Cunliffe, W. J. 1981. A review the microbiology of acne. *Journal of Applied Bacteriology* 51:195-215. 29.Holt, J. G., Krieg, N. R., Sneath, P. H. A., Staley, J. T. and Williams, S. T. 1994. Bergey ' s manual of determinative bacteriology. p. 580-596. Williams & Wilkins, Baltimore, MD, USA. 30.Ingham, E., Holland, K. T., Gowland, G. and Cunliffe, W. J. 1981. Partial purification and characterization of lipase (EC 3.1.1.3) from Propionibacterium acnes. *Journal of Microbiology* 124: 393-401. 31.Karen, W. M. and Edzard E. 2003. Herbal medicines for treatment of bacterial infections: a review of controlled clinical trials. *Journal of Antimicrobial Chemotherapy* 51:241-246 32.Katsambas, A. D., Stefanaki, C. and Cunliffe W. J. 2004. Guidelines for treating acne. *Clinics in Dermatology* 22:439-444. 33.Kolodney, M. S. and Marvi I. 2005. Acne fulminans with synovitis-acne-pustulosishyperostosis-osteitis (SAPHO) syndrome treated with infliximab. *Journal of the American Academy of Dermatology* 52:118-120. 34.Krautheim, A. and Gollnick, H. P. 2004. Acne: topical treatment. *Clinical Dermatology* 22:398-407. 35.Liao, S. 2001. The medicinal action of androgend and green tea epigallocatechin gallate. *Hong Kong Medical Journa*. 7: 369-374. 36.Lim, Y. H., Kim, I. H. and Seo, J. J. 2007. In vitro activity of kaempferol isolate from the Impatiens balsamina alone and in combination with erythromycin or clindamycin against Propionibacterium acnes. *The Journal of Microbiology*. 45: 473-477. 37.Lindor, N. M., Arsenault, M. T., Solomon, H., Seidman, E. C. and Mcevoy, T. M. 1997. A new autosomal dominant disorder of pyogenic sterile arthritis, pyoderma gangrenosum, and acne: PAPA syndrome. *Mayo Clinic Proceedings* 72:611-615. 38.Ljubojevic, S., Skerlev, M., Lipozencic, J. and Basta, J. A. 2002. The Role of Malassezia furfur in Dermatology. *Clinics in Dermatology* 20: 179-182. 39.Nam, C., Kim, S., Sim, Y. and Chang, I. 2003. Anti-acne effects of oriental herb extracts: anovel screening method to select anti-acne agents. *Skin Pharmacology and Applied Skin Physiology* 16: 84-90. 40.Orafidiya, L. O., Agbani, E. O., Oyedele, A. O., Babalola, O. O. and Onayemi, O. 2002. Preliminary clinical tests on topical preparations of Ocimum gratissimum Linn leaf essential oil for the treatment of acne vulgaris. *Clinical Drug Investigations* 22: 313-319. 41.Raman, A., Weir, U. and Bloomfield, S. F. 1995. Antimicrobial effects of tea-tree oil and its major components on Staphylococcus aureus, Staph. Epidermidis and Propionibacterium acnes. 1995. *Letters in Applied Microbiology* 21:242-245. 42.Ross, J. I., Snelling, A. M., Eady, E. A., Cove, J. H., Cunliffe, W. J., Leyden, J. J., Collignon, P., Dreno, B., Reynaud, A., Fluhr, J. and Oshima, S. 2001. Phenotypic and genotypic characterization of antibiotic-resistant Propionibacterium acnes isolated from acne patients attending dermatology clinics in Europe, the USA, Japan, and Australia. *British Journal for Dermatology* 144: 339-346. 43.Wang, L., Tu, Y. C., Lian, T. W., Hung, J. T., Yen, J. H., and Wu, M. J. 2006. Distinctive anti-inflammatory effects of flavonols. *Journal of Agricultural and Food Chemistry* 54: 9798-9804. 44.Webster, G. F. 2002. Acne vulgaris. *British Medical Journal* 325:475-479. 45.White, G. M. 1999. Acne therapy. *Disease-a-Month* 45:305-328. 46.Yang, M. H. and Choong, Y. M. 2001. A rapid gas chromatographic method for direct determination of short-chain (C2-C12) volatile organic acids in foods. *Food chemistry* 75:101-108. 47.Yang, X., Summerhurst, D. K., Koval, S. F., Ficker, M. L., Smith, M. L., and Bernards, M. A. 2001. Isolation of an antimicrobial compound from Impatiens balsamina L. using bioassay-guided fraction. *Phytotherapy Research* 15: 676-680